

t26_valued_1 (TMUD- pjP9QiSyti5mFWYgQoFRpTh4c8izRqT)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k61_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (r1_tarski (k10_xtuple_0 (k61_valued_1 X0 X1)) (k10_xtuple_0 X0))) \quad (1)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2.(v7_ordinal1 X2) \Rightarrow ((X2 \in k9_xtuple_0 X0) \Rightarrow (k2_xcmplx_0 X2 X1 \in k1_relset_1 k5_numbers (k61_valued_1 X0 X1)))))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (7)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge(v1_funct_1 X0))\wedge(v7_ordinal1 X1))\Rightarrow((v1_relat_1 (k61_valued_1 X0 X1))\wedge((v4_relat_1 (k61_valued_1 X0 X1) k5_numbers)\wedge(v1_funct_1 (k61_valued_1 X0 X1)))) \quad (10)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow(X2 \in X1)) \quad (12)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.(X1 = k10_xtuple_0 X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow(\exists X3.(X3 \in k9_xtuple_0 X0)\wedge(X2 = k1_funct_1 X0 X3)))) \quad (13)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.(v7_ordinal1 X1)\Rightarrow(\forall X2.((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((X2 = k61_valued_1 X0 X1)\Leftrightarrow((k9_xtuple_0 X2 = ReplSep (toset (\lambda X3 : \iota.m2_subset_1 X3 k1_numbers k5_numbers)) (\lambda X3 : \iota.X3 \in k9_xtuple_0 X0) (\lambda X3 : \iota.k2_nat_1 X3 X1))\wedge(\forall X3.(m2_subset_1 X3 k1_numbers k5_numbers)\Rightarrow((X3 \in k9_xtuple_0 X0)\Rightarrow(k1_funct_1 X2 (k2_nat_1 X3 X1) = k1_funct_1 X0 X3)))))))) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1)\Leftrightarrow((r1_tarski X0 X1)\wedge(r1_tarski X1 X0)) \quad (15)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (16)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((r1_tarski (k9_xtuple_0 X0) k5_numbers) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (k10_xtuple_0 (k61_valued_1 X0 X1) = k10_xtuple_0 X0)))$$