

t27_bvfunc_1

(TMTpw7Yxv9tAr5zryjtBUfrsaNDXZ4b7zbn)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k14_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k13_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k8_margrel1 : \iota$ be given. Let $k2_xboolean : \iota$ be given. Let $k7_margrel1 : \iota$ be given. Let $k1_xboolean : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_bvfunc_1 : \iota \Rightarrow \iota$ be given. Let $k11_bvfunc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboolean : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ & X0) \Rightarrow (r1_xxreal_0 (k3_funct_2 X0 k6_margrel1 (k13_bvfunc_1 X0 \\ & X1) X2) (k3_funct_2 X0 k6_margrel1 X1 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$r1_xxreal_0 np_1 np_1 \quad (2)$$

Assume the following.

$$k8_margrel1 = k2_xboolean \quad (3)$$

Assume the following.

$$k7_margrel1 = k1_xboolean \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((\neg v1_xboole_0 X0) \wedge \\ & (((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1)))))) \wedge (m1_subset_1 X3 X0))) \Rightarrow (k3_funct_2 X0 \\ & X1 X2 X3 = k1_funct_1 X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ & (((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))))))\wedge(m1_subset_1 X3 X0))\Rightarrow(m1_subset_1 (\\ & k3_funct_2 X0 X1 X2 X3) X1) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow((v1_funct_1 (k12_bvfunc_1 X0))\wedge \\ & ((v1_funct_2 (k12_bvfunc_1 X0) X0 k6_margrel1)\wedge(m1_subset_1 \\ & (k12_bvfunc_1 X0) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow((v1_funct_1 (k11_bvfunc_1 X0))\wedge \\ & ((v1_funct_2 (k11_bvfunc_1 X0) X0 k6_margrel1)\wedge(m1_subset_1 \\ & (k11_bvfunc_1 X0) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.(v1_xboolean X0)\Leftrightarrow((X0 = k1_xboolean)\vee(X0 = k2_xboolean)) \quad (9)$$

Assume the following.

$$k2_xboolean = np_1 \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(\\ & (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(((\forall X2.(m1_subset_1 \\ & X2 X0)\Rightarrow(k3_funct_2 X0 k6_margrel1 X1 X2 = k7_margrel1))\Rightarrow(k14_bvfunc_1 \\ & X0 X1 = k11_bvfunc_1 X0))\wedge((\neg\forall X2.(m1_subset_1 X2 X0)\Rightarrow(k3_funct_2 \\ & X0 k6_margrel1 X1 X2 = k7_margrel1))\Rightarrow(k14_bvfunc_1 X0 X1 = k12_bvfunc_1 \\ & X0)))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(\\ & (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(((\forall X2.(m1_subset_1 \\ & X2 X0)\Rightarrow(k3_funct_2 X0 k6_margrel1 X1 X2 = k8_margrel1))\Rightarrow(k13_bvfunc_1 \\ & X0 X1 = k12_bvfunc_1 X0))\wedge((\neg\forall X2.(m1_subset_1 X2 X0)\Rightarrow(k3_funct_2 \\ & X0 k6_margrel1 X1 X2 = k8_margrel1))\Rightarrow(k13_bvfunc_1 X0 X1 = k11_bvfunc_1 \\ & X0)))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((X1 = k12_bvfunc_1 X0) \Leftrightarrow (\forall X2. \\ (m1_subset_1 X2 X0) \Rightarrow (k3_funct_2 X0 k6_margrel1 X1 X2 = k8_margrel1)))) \end{aligned} \quad (13)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((X1 = k11_bvfunc_1 X0) \Leftrightarrow (\forall X2. \\ (m1_subset_1 X2 X0) \Rightarrow (k3_funct_2 X0 k6_margrel1 X1 X2 = k7_margrel1)))) \end{aligned} \quad (14)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k6_margrel1) \Rightarrow (v1_xboolean X0) \quad (15)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.(m1_subset_1 X2 \\ X0) \Rightarrow (r1_xxreal_0 (k3_funct_2 X0 k6_margrel1 X1 X2) (k3_funct_2 \\ X0 k6_margrel1 (k14_bvfunc_1 X0 X1) X2)))) \end{aligned}$$