

## t27\_bvfunc\_7

(TMSz2nvire7jPVpmU64hQ2kahqBadxxGLTC)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k12\_bvfunc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r1\_bvfunc\_1 X0 X1 (k10\_bvfunc\_1 \\ & X0 (k10\_bvfunc\_1 X0 (k10\_bvfunc\_1 X0 X1 X2) (k10\_bvfunc\_1 X0 X2 X1)) \\ & X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow ((r2\_funct\_2 X0 k6\_margrel1 \\ & (k10\_bvfunc\_1 X0 X1 X2) (k12\_bvfunc\_1 X0)) \Leftrightarrow (r2\_funct\_2 X0 k6\_margrel1 \\ & X1 X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1)))))) \wedge ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 \\ & X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \Rightarrow ((r2\_funct\_2 X0 X1 X2 \\ & X3) \Leftrightarrow (X2 = X3)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(((v1\_funct\_1 \\ & X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1))))))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 \\ & X2 X0 k6\_margrel1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 k6\_margrel1))))))\Rightarrow((v1\_funct\_1 (k2\_bfunc\_1 X0 X1 X2))\wedge( \\ & (v1\_funct\_2 (k2\_bfunc\_1 X0 X1 X2) X0 k6\_margrel1)\wedge(m1\_subset\_1 \\ & (k2\_bfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow((v1\_funct\_1 (k12\_bfunc\_1 X0))\wedge \\ & ((v1\_funct\_2 (k12\_bfunc\_1 X0) X0 k6\_margrel1)\wedge(m1\_subset\_1 \\ & (k12\_bfunc\_1 X0) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(((v1\_funct\_1 \\ & X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1))))))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 \\ & X2 X0 k6\_margrel1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 k6\_margrel1))))))\Rightarrow((v1\_funct\_1 (k10\_bfunc\_1 X0 X1 X2))\wedge \\ & ((v1\_funct\_2 (k10\_bfunc\_1 X0 X1 X2) X0 k6\_margrel1)\wedge(m1\_subset\_1 \\ & (k10\_bfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(((v1\_funct\_1 \\ & X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1))))))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 \\ & X2 X0 k6\_margrel1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 k6\_margrel1))))))\Rightarrow(k2\_bfunc\_1 X0 X1 X2 = k2\_bfunc\_1 X0 X2 \\ & X1) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0)\wedge(((v1\_funct\_1 \\ & X1)\wedge((v1\_funct\_2 X1 X0 k6\_margrel1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1))))))\wedge((v1\_funct\_1 X2)\wedge((v1\_funct\_2 \\ & X2 X0 k6\_margrel1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 k6\_margrel1))))))\Rightarrow(k10\_bfunc\_1 X0 X1 X2 = k10\_bfunc\_1 X0 \\ & X2 X1) \end{aligned} \quad (8)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge \\ (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge \\ ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r1\_bvfunc\_1 X0 X1 (k10\_bvfunc\_1 \\ X0 (k10\_bvfunc\_1 X0 (k2\_bvfunc\_1 X0 X1 X2) (k2\_bvfunc\_1 X0 X2 X1)) \\ X1)))) \end{aligned}$$