

t27_finseq_4 (TMX- uWvRiCW9Ju68j2uhRrAb1VxwJ54SPLmw)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_finseq_4 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(r2_finseq_4 \\ & X0 X1) \Leftrightarrow (\exists X2.(X2 \in k9_xtuple_0 X0) \wedge ((X1 = k1_funct_1 X0 X2) \wedge \\ & (\forall X3. \neg (X3 \in k9_xtuple_0 X0) \wedge ((X3 \neq X2) \wedge (k1_funct_1 X0 X3 = \\ & X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(X1 \in k10_xtuple_0 X0) \Rightarrow (k4_finseq_4 X0 X1 \in k4_finseq_1 \\ & X0)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(X1 \in k10_xtuple_0 X0) \Rightarrow (k1_funct_1 X0 (k4_finseq_4 \\ & X0 X1) = X1)) \end{aligned} \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m1_subset_1 (k4_finseq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (8)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1.((X1 \in k10_xtuple_0 X0) \wedge (\forall X2.(v7_ordinal1 X2) \Rightarrow (\neg(X2 \in k4_finseq_1 X0) \wedge ((X2 \neq k4_finseq_4 X0 X1) \wedge (k1_funct_1 X0 X2 = X1))))) \Rightarrow (r2_finseq_4 X0 X1))$$