

t27_funct_4

(TMJLUS9bZKTzU1s2UwDv8RTZtxJQy1ZvQAL)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_partfun1 X0 X1) \Leftrightarrow (\exists X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \wedge ((r1_tarski X0 X2) \wedge (r1_tarski X1 X2))))) \quad (1)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (r1_tarski X0 (k1_funct_4 X1 X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k1_funct_4 X0 X1)) \wedge (v1_funct_1 (k1_funct_4 X0 X1))) \quad (4)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((r1_partfun1 X0 (k1_funct_4 X1 X2)) \Rightarrow (r1_partfun1 X0 X2))))$$