

t27_group_1
(TMW6J3hqtQVPHdQSrJTnNFYkFLJBVtBZz5o)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_group_1 : \iota \Rightarrow o$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_group_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k6_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (k5_group_1 X0 np_1 X1 = X1)) \quad (1)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (2)$$

Assume the following.

$$k2_xcmplx_0 np_1 np_1 = np_2 \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v7_ordinal1 X0) \wedge (m1_subset_1 X1 k5_numbers)) \Rightarrow (k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. ((\neg v2_struct_0 X1) \wedge ((v2_group_1 X1) \wedge ((v3_group_1 X1) \wedge (l3_algstr_0 X1)))) \Rightarrow (\forall X2. (m1_subset_1 X2 (u1_struct_0 X1)) \Rightarrow (k5_group_1 X1 (k1_nat_1 X0 np_1) X2 = k6_algstr_0 X1 (k5_group_1 X1 X0 X2) X2))) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Theorem 1

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 \\ X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 \\ X0)) \Rightarrow (k5_group_1 X0 \text{ np_2 } X1 = k6_algstr_0 X0 X1 X1)) \end{aligned}$$