

t27\_int\_2 (TMMxd-  
pUk4rTwKzYe8zN25D7yugKm7WXHuaG)

October 27, 2020

Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Let  $r1\_int\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_int\_1 X0) \Rightarrow ((r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (X0 \in k5\_numbers)) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2.(v1\_xxreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 X0 X2)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (3)$$

Assume the following.

$$m1\_subset\_1 k1\_xboole\_0 k4\_ordinal1 \quad (4)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (5)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((r1\_int\_1 X1 X0) \Rightarrow ((r1\_xxreal\_0 X0 k6\_numbers) \vee (r1\_xxreal\_0 X1 X0)))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0)\wedge(v1\_xxreal\_0 X1))\Rightarrow( (r1\_xxreal\_0 X0 X1)\vee(r1\_xxreal\_0 X1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow(v1\_xxreal\_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1\_int\_1 X0)\Rightarrow(v1\_xreal\_0 X0) \quad (11)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(v1\_xxreal\_0 X0) \quad (12)$$

**Theorem 1**

$$\forall X0.(v1\_int\_1 X0)\Rightarrow(\forall X1.(v1\_int\_1 X1)\Rightarrow((r1\_int\_1 X1 X0)\Rightarrow((r1\_xxreal\_0 X0 k6\_numbers)\vee(r1\_xxreal\_0 X1 X0))))$$