

t27\_mboolean  
(TMJnDp1pXopizcW9TztdzfmH8hgKp9ovfH3)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r6\_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_mboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_pboole : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_tarski : \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. k3\_tarski (k2\_xboole\_0 X0 X1) = k2\_xboole\_0 (k3\_tarski X0) (k3\_tarski X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 \\ & X1 X0) \wedge ((v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \wedge ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))))) \Rightarrow \\ & (r6\_pboole X0 X1 X2) \Leftrightarrow (X1 = X2) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0))) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow \\ & (\forall X3. (X3 \in X0) \Rightarrow (k1\_funct\_1 (k2\_mboolean X0 (k2\_pboole X0 \\ & X1 X2)) X3 = k3\_tarski (k2\_xboole\_0 (k1\_funct\_1 X1 X3) (k1\_funct\_1 \\ & X2 X3)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 \\ & X1 X0) \wedge ((v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \wedge ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))))) \Rightarrow \\ & (k2\_pboole X0 X1 X1 = X1) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 \\ & X1 X0) \wedge ((v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \wedge ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))))) \Rightarrow \\ & ((v1\_relat\_1 (k2\_pboole X0 X1 X2)) \wedge ((v4\_relat\_1 (k2\_pboole X0 \\ & X1 X2) X0) \wedge ((v1\_funct\_1 (k2\_pboole X0 X1 X2)) \wedge (v1\_partfun1 (k2\_pboole \\ & X0 X1 X2) X0)))) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge ( \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \Rightarrow ((v1\_relat\_1 (k2\_mboolean \\ & X0 X1)) \wedge ((v4\_relat\_1 (k2\_mboolean X0 X1) X0) \wedge ((v1\_funct\_1 (k2\_mboolean \\ & X0 X1)) \wedge (v1\_partfun1 (k2\_mboolean X0 X1) X0)))) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge ( \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow \\ & (\forall X3. ((v1\_relat\_1 X3) \wedge ((v4\_relat\_1 X3 X0) \wedge ((v1\_funct\_1 \\ & X3) \wedge (v1\_partfun1 X3 X0)))) \Rightarrow ((X3 = k2\_pboole X0 X1 X2) \Leftrightarrow (\forall X4. \\ & (X4 \in X0) \Rightarrow (k1\_funct\_1 X3 X4 = k2\_xboole.0 (k1\_funct\_1 X1 X4) (k1\_funct\_1 \\ & X2 X4)))))) \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge ( \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow \\ & ((X2 = k2\_mboolean X0 X1) \Leftrightarrow (\forall X3. (X3 \in X0) \Rightarrow (k1\_funct\_1 X2 X3 = \\ & k3\_tarski (k1\_funct\_1 X1 X3)))))) \end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 \\ & X1 X0) \wedge ((v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \wedge ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))))) \Rightarrow \\ & (k2\_pboole X0 X1 X2 = k2\_pboole X0 X2 X1) \end{aligned} \tag{9}$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge ( \\ & (v1\_funct\_1 X1) \wedge (v1\_partfun1 X1 X0)))) \Rightarrow (\forall X2. ((v1\_relat\_1 \\ & X2) \wedge ((v4\_relat\_1 X2 X0) \wedge ((v1\_funct\_1 X2) \wedge (v1\_partfun1 X2 X0)))) \Rightarrow \\ & (r6\_pboole X0 (k2\_mboolean X0 (k2\_pboole X0 X1 X2)) (k2\_pboole X0 \\ & (k2\_mboolean X0 X1) (k2\_mboolean X0 X2)))) \end{aligned}$$