

t27_seqfunc
(TMXQ4sdCTxSVaB4Wf3HtJHQzjskWbdN92ji)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_seqfunc : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_seqfunc : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_seqfunc : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 k5_numbers (k4_partfun1 X0 k1_numbers)) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k4_partfun1 X0 k1_numbers)))))) \Rightarrow \\ & (\forall X2.(r1_seqfunc X0 k1_numbers X1 X2) \Rightarrow (\forall X3.(m1_subset_1 \\ & X3 X0) \Rightarrow ((X3 \in X2) \Rightarrow (r1_seqfunc X0 k1_numbers X1 (k1_tarski X3)))))) \\ & \hspace{15em} (1) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 k5_numbers (k4_partfun1 X0 k1_numbers)) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k4_partfun1 X0 k1_numbers)))))) \Rightarrow \\ & (\forall X2.(r2_seqfunc X0 X1 X2) \Leftrightarrow ((r1_seqfunc X0 k1_numbers X1 \\ & X2) \wedge (\exists X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k1_numbers)))))) \wedge ((X2 = k1_relset_1 X0 X3) \wedge (\forall X4. \\ & (m1_subset_1 X4 X0) \Rightarrow ((X4 \in X2) \Rightarrow (\forall X5.(m1_subset_1 X5 k1_numbers) \Rightarrow \\ & (\neg(\neg r1_xxreal_0 X5 k6_numbers) \wedge (\forall X6.(m2_subset_1 X6 k1_numbers \\ & k5_numbers) \Rightarrow (\exists X7.(m2_subset_1 X7 k1_numbers k5_numbers) \wedge \\ & ((r1_xxreal_0 X6 X7) \wedge (r1_xxreal_0 X5 (k18_complex1 (k9_real_1 \\ & (k1_seq_1 (k1_seqfunc X0 k1_numbers X1 X7) X4) (k1_seq_1 X3 X4)))))))))))))) \\ & \hspace{15em} (2) \end{aligned}$$

Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 k5_numbers (k4_partfun1 X0 k1_numbers)) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k4_partfun1 X0 k1_numbers)))))) \Rightarrow \\ & (\forall X2. (r2_seqfunc X0 X1 X2) \Rightarrow (\forall X3. (m1_subset_1 X3 \\ & X0) \Rightarrow ((X3 \in X2) \Rightarrow (r1_seqfunc X0 k1_numbers X1 (k1_tarSKI X3)))))) \end{aligned}$$