

## t27\_waybel30

(TMdjB1ZBaoCRC7NsiG3XuEojcvTqQWKMKsw)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v3\_orders\_2 : \iota \Rightarrow o$  be given. Let  $l1\_orders\_2 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_waybel30 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. r1\_tarski X0 (k2\_xboole\_0 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 (k1\_zfmisc\_1 X1)) \Leftrightarrow (r1\_tarski X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v3\_orders\_2 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \Rightarrow ((r1\_tarski X1 X2) \Rightarrow (r1\_tarski (k1\_waybel30 X0 X1) (k1\_waybel30 X0 X2)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0))) \Rightarrow (k4\_subset\_1 X0 X1 X2 = k2\_xboole\_0 X1 X2) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0))) \Rightarrow (m1\_subset\_1 (k4\_subset\_1 X0 X1 X2) (k1\_zfmisc\_1 X0)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(((\neg v2\_struct\_0 X0)\wedge((v3\_orders\_2 X0)\wedge(l1\_orders\_2 X0)))\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0))))\Rightarrow(m1\_subset\_1 (k1\_waybel30 X0 X1) (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.k2\_xboole\_0 X0 X1 = k2\_xboole\_0 X1 X0 \quad (7)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0)\wedge((v3\_orders\_2 X0)\wedge(l1\_orders\_2 X0)))\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 X0)))\Rightarrow(\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (u1\_struct\_0 X0)))\Rightarrow(r1\_tarski (k4\_subset\_1 (u1\_struct\_0 X0) (k1\_waybel30 X0 X1) (k1\_waybel30 X0 X2)) (k1\_waybel30 X0 (k4\_subset\_1 (u1\_struct\_0 X0) X1 X2)))))) \end{aligned}$$