

t27\_xxreal\_0

(TMLV92uWhJCVggKdFaKkKjrXmLU8MWeeBzg)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow \\ & (r1\_xxreal\_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (( \\ & (r1\_xxreal\_0 X1 X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X0)) \wedge ((\neg r1\_xxreal\_0 X1 \\ & X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow ( \\ & (r1\_xxreal\_0 X0 X1) \vee (r1\_xxreal\_0 X1 X0)) \end{aligned} \quad (3)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (\forall X3.(v1\_xxreal\_0 X3) \Rightarrow (\neg(\neg r1\_xxreal\_0 \\ & X1 X0) \wedge ((\neg r1\_xxreal\_0 X3 X2) \wedge (r1\_xxreal\_0 (k4\_xxreal\_0 X1 X3) \\ & (k4\_xxreal\_0 X0 X2)))))))) \end{aligned}$$