

t27_zf_lang1

(TMGpchkP8nZbRuScoPvzUuPSjGAsagQrubG)

October 27, 2020

Let $v1_zf_lang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v10_zf_lang : \iota \Rightarrow o$ be given. Let $k27_zf_lang : \iota \Rightarrow \iota$ be given. Let $k25_zf_lang : \iota \Rightarrow \iota$ be given. Let $k21_zf_lang : \iota \Rightarrow \iota$ be given. Let $k26_zf_lang : \iota \Rightarrow \iota$ be given. Let $k22_zf_lang : \iota \Rightarrow \iota$ be given. Let $k12_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k28_zf_lang : \iota \Rightarrow \iota$ be given. Let $k11_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow ((k27_zf_lang \\ & (k12_zf_lang X0 X1) = X0) \wedge (k28_zf_lang (k12_zf_lang X0 X1) = X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow ((k25_zf_lang \\ & (k11_zf_lang X0 X1) = X0) \wedge (k26_zf_lang (k11_zf_lang X0 X1) = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow ((k21_zf_lang \\ & (k7_zf_lang X0 X1) = X0) \wedge (k22_zf_lang (k7_zf_lang X0 X1) = X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow ((\\ & v10_zf_lang X0) \Rightarrow (X0 = k12_zf_lang (k27_zf_lang X0) (k28_zf_lang \\ & X0))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_zf_lang X0) \wedge (m1_finseq_1 X0 k5_numbers)) \wedge \\ & ((v1_zf_lang X1) \wedge (m1_finseq_1 X1 k5_numbers)) \Rightarrow ((v1_zf_lang \\ & (k11_zf_lang X0 X1)) \wedge (m2_finseq_1 (k11_zf_lang X0 X1) k5_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow ((\\ & v10_zf_lang X0) \Leftrightarrow (\exists X1. ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 \\ & k5_numbers)) \wedge (\exists X2. ((v1_zf_lang X2) \wedge (m2_finseq_1 X2 k5_numbers)) \wedge \\ & (X0 = k12_zf_lang X1 X2)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow (k12_zf_lang \\ & X0 X1 = k7_zf_lang (k11_zf_lang X0 X1) (k11_zf_lang X1 X0))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow ((\\ & v10_zf_lang X0) \Rightarrow ((k27_zf_lang X0 = k25_zf_lang (k21_zf_lang X0)) \wedge \\ & (k27_zf_lang X0 = k26_zf_lang (k22_zf_lang X0)))) \end{aligned}$$