

t28_arytm_1

(TMbMbG2KGZ17sYBd3v1RCS6XJ9w9uagPbGv)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_arytm_2 : \iota$ be given. Let $r1_arytm_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_arytm_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_arytm_3 : \iota$ be given. Let $k2_arytm_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_arytm_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k2_arytm_2) \Rightarrow (\forall X2.(m1_subset_1 X2 k2_arytm_2) \Rightarrow (\neg(\neg \\ & r1_arytm_2 X0 X1) \wedge ((X2 \neq k11_arytm_3) \wedge (k4_tarski k11_arytm_3 \\ & (k8_arytm_2 X2 (k1_arytm_1 X0 X1)) \neq k2_arytm_1 (k8_arytm_2 X2 X1) \\ & (k8_arytm_2 X2 X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k2_arytm_2) \Rightarrow (((r1_arytm_2 X0 X1) \wedge (r1_arytm_2 X1 X0)) \Rightarrow (X0 = \\ & X1))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (k1_arytm_1 X0 X0 = k11_arytm_3) \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k2_arytm_2) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k2_arytm_2) \Rightarrow (\forall X2.(m1_subset_1 X2 k2_arytm_2) \Rightarrow ((r1_arytm_2 \\ & X1 X0) \Rightarrow ((k1_arytm_1 X0 X1 = k11_arytm_3) \vee ((X2 = k11_arytm_3) \vee \\ & k2_arytm_1 (k8_arytm_2 X2 X1) (k8_arytm_2 X2 X0) = k4_tarski k11_arytm_3 \\ & (k8_arytm_2 X2 (k1_arytm_1 X0 X1)))))) \end{aligned}$$