

t28_card_1
(TMcBkGzEsDgGwtaqfNK5ugcsc6TVEQiWtif)

October 27, 2020

Let $r2_wellord2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X1 \in X0) \Rightarrow (k1_funct_1 (k2_funcop_1 X0 X2) X1 = X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((k9_xtuple_0 X1 = k1_tarski X0) \Rightarrow (k10_xtuple_0 X1 = k1_tarski (k1_funct_1 X1 X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r2_wellord2 X0 X1) \Rightarrow (r2_wellord2 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k2_funcop_1 X0 X1)) \wedge (v1_funct_1 (k2_funcop_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (r2_wellord2 X0 X1) \Leftrightarrow (\exists X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \wedge ((v2_funct_1 X2) \wedge ((k9_xtuple_0 X2 = X0) \wedge (k10_xtuple_0 X2 = X1)))) \quad (6)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v2_funct_1 X0) \Leftrightarrow (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \wedge ((X2 \in k9_xtuple_0 X0) \wedge (k1_funct_1 X0 X1 = k1_funct_1 X0 X2)))) \Rightarrow (X1 = X2))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarSKI X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (8)$$

Theorem 1

$$\forall X0.\forall X1.(r2_wellord2 X0 (k1_tarSKI X1)) \Leftrightarrow (\exists X2. X0 = k1_tarSKI X2)$$