

## t28\_euclid

(TMK36MKwykQkfgViWSH1wzdnqTLCeKSegyR)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k1\_rlvect\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $k4\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v2\_pre\_topc : \iota \Rightarrow o$  be given. Let  $v5\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l2\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $l2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $l1\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge \\ (v13\_algstr\_0 X1) \wedge ((v2\_rlvect\_1 X1) \wedge ((v3\_rlvect\_1 X1) \wedge ((v4\_rlvect\_1 \\ X1) \wedge ((v5\_rlvect\_1 X1) \wedge ((v6\_rlvect\_1 X1) \wedge ((v7\_rlvect\_1 X1) \wedge \\ ((v8\_rlvect\_1 X1) \wedge (l1\_rlvect\_1 X1)))))))))) \Rightarrow (\forall X2.(m1\_subset\_1 \\ X2 (u1\_struct\_0 X1)) \Rightarrow (((X0 = k6\_numbers) \vee (X2 = k4\_struct\_0 X1)) \Rightarrow \\ (k1\_rlvect\_1 X1 X2 X0 = k4\_struct\_0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(v7\_ordinal1 X0) \Rightarrow ((v2\_pre\_topc (k15\_euclid X0)) \wedge \\ ((v13\_algstr\_0 (k15\_euclid X0)) \wedge ((v2\_rlvect\_1 (k15\_euclid X0)) \wedge \\ ((v3\_rlvect\_1 (k15\_euclid X0)) \wedge ((v4\_rlvect\_1 (k15\_euclid X0)) \wedge \\ ((v5\_rlvect\_1 (k15\_euclid X0)) \wedge ((v6\_rlvect\_1 (k15\_euclid X0)) \wedge \\ ((v7\_rlvect\_1 (k15\_euclid X0)) \wedge ((v8\_rlvect\_1 (k15\_euclid X0)) \wedge \\ (v5\_rltopsp1 (k15\_euclid X0)))))))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((\neg v2\_struct\_0 (k15\_euclid X0)) \wedge \\ (v5\_rltopsp1 (k15\_euclid X0))) \tag{3}$$

Assume the following.

$$\forall X0.(l2\_algstr\_0 X0) \Rightarrow ((l2\_struct\_0 X0) \wedge (l1\_algstr\_0 X0)) \tag{4}$$

Assume the following.

$$\forall X0.(l1\_rlvect\_1 X0) \Rightarrow (l2\_algstr\_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.(l1\_rltopsp1 X0) \Rightarrow ((l1\_rlvect\_1 X0) \wedge (l1\_pre\_topc X0)) \quad (6)$$

Assume the following.

$$\forall X0.(l2\_struct\_0 X0) \Rightarrow (m1\_subset\_1 (k4\_struct\_0 X0) (u1\_struct\_0 X0)) \quad (7)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((v5\_rltopsp1 (k15\_euclid X0)) \wedge (l1\_rltopsp1 (k15\_euclid X0))) \quad (8)$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (k1\_rlvect\_1 (k15\_euclid X0) (k4\_struct\_0 (k15\_euclid X0)) X1 = k4\_struct\_0 (k15\_euclid X0)))$$