

t28_finseq_6
(TMQVjyJfQ2E4qX4Ak3znukXV7aWZefbiTST)

October 27, 2020

Let $k5_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $np_3 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. k5_relat_1 (k11_finseq_1 X0 X1 X2) (k2_finseq_1 np_2) = k10_finseq_1 X0 X1 \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1. \forall X2. (v7_ordinal1 X2) \Rightarrow (((X1 \in k10_xtuple_0 X0) \wedge (X2 = k6_xcmplx_0 (k4_finseq_4 X0 X1) np_1)) \Rightarrow (k5_relat_1 X0 (k2_finseq_1 X2) = k5_finseq_4 X0 X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \neq X1) \wedge ((X2 \neq X1) \wedge (k4_finseq_4 (k11_finseq_1 X0 X2 X1) X1 \neq np_3)) \quad (3)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k6_xcmplx_0 np_3 np_1 = np_2 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.k10_xtuple_0 (k11_finseq_1 X0 X1 X2) = k1_enumset1 X0 X1 X2 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(v1_relat_1 (k11_finseq_1 X0 X1 X2)) \wedge (v1_funct_1 (k11_finseq_1 X0 X1 X2)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.v1_finseq_1 (k11_finseq_1 X0 X1 X2) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(X3 = k1_enumset1 X0 X1 X2) \Leftrightarrow (\forall X4.(X4 \in X3) \Leftrightarrow (\neg(X4 \neq X0) \wedge ((X4 \neq X1) \wedge (X4 \neq X2)))) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (11)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.\neg(X0 \neq X1) \wedge ((X2 \neq X1) \wedge (k5_finseq_4 (k11_finseq_1 X0 X2 X1) X1 \neq k10_finseq_1 X0 X2))$$