

t28\_hilbert1  
(TMPfxGHbicS4L7qThU7btpDtjpkavsrDVUg)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_hilbert1 : \iota$  be given. Let  $k3\_hilbert1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_hilbert1 : \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_subset\_1 X0 k1\_hilbert1) \Rightarrow (\forall X1.(m1\_subset\_1 \\ X1 k1\_hilbert1) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_hilbert1) \Rightarrow (( \\ k3\_hilbert1 X0 (k3\_hilbert1 X1 X2) \in k6\_hilbert1) \Rightarrow (k3\_hilbert1 \\ X1 (k3\_hilbert1 X0 X2) \in k6\_hilbert1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_hilbert1) \Rightarrow (k3\_hilbert1 X0 X0 \in k6\_hilbert1) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_hilbert1) \wedge (m1\_subset\_1 \\ X1 k1\_hilbert1)) \Rightarrow (m1\_subset\_1 (k3\_hilbert1 X0 X1) k1\_hilbert1) \quad (3)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(m1\_subset\_1 X0 k1\_hilbert1) \Rightarrow (\forall X1.(m1\_subset\_1 \\ X1 k1\_hilbert1) \Rightarrow (k3\_hilbert1 X0 (k3\_hilbert1 (k3\_hilbert1 X0 \\ X1) X1) \in k6\_hilbert1)) \end{aligned}$$