

t28_hilbert2 (TMTE-
mEUmkcNQ4st1KCzk7ZLywpvYZRrKbYw)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_hilbert1 : \iota$ be given. Let $k3_hilbert1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v4_funct_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_hilbert1) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 k1_hilbert1) \Rightarrow ((\neg r1_xreal_0 (k3_finseq_1 (k3_hilbert1 X0 \\ & X1)) (k3_finseq_1 X0)) \wedge (\neg r1_xreal_0 (k3_finseq_1 (k3_hilbert1 \\ & X0 X1)) (k3_finseq_1 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (2)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (3)$$

Assume the following.

$$v4_funct_1 k1_hilbert1 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Rightarrow (m1_subset_1 X2 X0)) \end{aligned} \quad (5)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow (r1_xreal_0 X0 X1)\vee(r1_xreal_0 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(v4_funct_1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1))) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xreal_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_hilbert1)\Rightarrow(v1_finseq_1 X0) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_xboole_0 X1)) \quad (13)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k1_hilbert1)\Rightarrow(\forall X1.(m1_subset_1 X1 k1_hilbert1)\Rightarrow((k3_hilbert1 X0 X1\neq X0)\wedge(k3_hilbert1 X0 X1\neq X1)))$$