

t28_lpspace1 (TMLvct- NCJH8CRpVDRAP4oQUtEjcfdjkaCB4)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r1_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r3_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow ((r2_relset_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (2)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \wedge (m1_subset_1 X3 X0))) \Rightarrow (k3_funct_2 X0 X1 X2 X3 = k1_funct_1 X2 X3) \quad (4)$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (((v1_funct_1 (k8_lpspace1 X0 k6_numbers)) \wedge (m1_subset_1 (k8_lpspace1 \\
& X0 k6_numbers) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \wedge (\\
& k8_lpspace1 X0 k6_numbers \in k9_lpspace1 X0 X1 X2)))
\end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2) \wedge \\
& (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow ((v1_funct_1 \\
& (k2_partfun1 X0 X1 X2 X3)) \wedge (m1_subset_1 (k2_partfun1 X0 X1 X2 X3) \\
& (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))
\end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (k9_lpspace1 X0 X1 X2 = ReplSep (toset (\lambda X3 : \iota.(v1_funct_1 \\
& X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \\
& (\lambda X3 : \iota.\exists X4.(m2_subset_1 X4 (k1_zfmisc_1 X0) X1) \wedge \\
& ((k3_funct_2 X1 k7_numbers X2 X4 = k6_numbers) \wedge ((k1_reset_1 X0 \\
& X3 = k3_subset_1 X0 X4) \wedge (r3_mesfunc6 X0 X1 X2 X3)))) (\lambda X3 : \iota. \\
& X3)))
\end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k1_numbers)))) \Rightarrow (\forall X4.((v1_funct_1 X4) \wedge (m1_subset_1 \\
& X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow ((r1_lpspace1 \\
& X0 X1 X2 X3 X4) \Leftrightarrow (\exists X5.(m2_subset_1 X5 (k1_zfmisc_1 X0) X1) \wedge \\
& ((k3_funct_2 X1 k7_numbers X2 X5 = k6_numbers) \wedge (r2_reset_1 X0 \\
& k1_numbers (k2_partfun1 X0 k1_numbers X3 (k3_subset_1 X0 X5)) (\\
& k2_partfun1 X0 k1_numbers X4 (k3_subset_1 X0 X5))))))
\end{aligned} \tag{8}$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\ & X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\ & (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k1_numbers)))) \Rightarrow (r1_lpspace1 X0 X1 X2 X3 X3)))) \end{aligned}$$