

t28_matroid0
(TMcw1X7GeUVjZCVcPcyeLMc5VWcHgtoG85v)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v3_pencil_1 : \iota \Rightarrow o$ be given. Let $v1_matroid0 : \iota \Rightarrow o$ be given. Let $v2_matroid0 : \iota \Rightarrow o$ be given. Let $v4_matroid0 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r2_matroid0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_matroid0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\neg(X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Rightarrow (k2_xboole_0 X0 X1 = X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (k4_subset_1 X0 X1 X2 = k2_xboole_0 X1 X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (m1_subset_1 (k6_domain_1 X0 X1) (k1_zfmisc_1 X0)) \quad (6)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((\neg v3_pencil_1 X0) \wedge ((v1_matroid0 \\
& X0) \wedge ((v2_matroid0 X0) \wedge ((v4_matroid0 X0) \wedge (l1_pre_topc X0)))))) \Rightarrow \\
& (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 \\
& X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((r2_matroid0 X0 X1 X2) \Leftrightarrow (k4_matroid0 \\
& X0 (k4_subset_1 (u1_struct_0 X0) X2 (k6_domain_1 (u1_struct_0 \\
& X0) X1)) = k4_matroid0 X0 X2))))))
\end{aligned} \tag{7}$$

Assume the following.

$$\forall X0. \forall X1. k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \tag{8}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((\neg v3_pencil_1 X0) \wedge ((v1_matroid0 \\
& X0) \wedge ((v2_matroid0 X0) \wedge ((v4_matroid0 X0) \wedge (l1_pre_topc X0)))))) \Rightarrow \\
& (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow \\
& (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow ((X2 \in X1) \Rightarrow (r2_matroid0 \\
& X0 X2 X1))))))
\end{aligned}$$