

t28_mesfunc6 (TMKcFGDFdRrL- RTEVVV5HCoGSdZMs5gym4PA)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $k7_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_mesfunc5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k26_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k32_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_mesfunc1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $k30_valued_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k24_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \Rightarrow (\\ r2_relset_1 X0 k7_numbers (k7_mesfunc1 X0 X1) (k6_mesfunc1 X0 X1 \\ (k1_real_1 np_1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ \forall X2.(m1_subset_1 X2 k1_numbers) \Rightarrow (r2_relset_1 X0 k7_numbers \\ (k1_mesfunc5 X0 (k26_valued_1 X0 k1_numbers X1 X2)) (k6_mesfunc1 \\ X0 (k1_mesfunc5 X0 X1) X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((r2_relset_1 X0 X1 X2 X3)\Leftrightarrow(X2 = X3)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v3_membered X1)\wedge((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))\Rightarrow(k32_valued_1 X0 X1 X2 = k30_valued_1 X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v3_membered X1)\wedge(((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge(v1_xreal_0 X3)))\Rightarrow(k26_valued_1 X0 X1 X2 X3 = k24_valued_1 X2 X3) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(k1_real_1 X0 = k4_xcmplx_0 X0) \quad (7)$$

Assume the following.

$$v3_membered k1_numbers \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_relat_1 X1)\wedge((v4_relat_1 X1 X0)\wedge((v5_relat_1 X1 k7_numbers)\wedge(v1_funct_1 X1))))\Rightarrow((v1_funct_1 (k7_mesfunc1 X0 X1))\wedge(m1_subset_1 (k7_mesfunc1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_relat_1 X1)\wedge((v4_relat_1 X1 X0)\wedge((v5_relat_1 X1 k7_numbers)\wedge(v1_funct_1 X1))))\wedge(m1_subset_1 X2 k1_numbers)))\Rightarrow((v1_funct_1 (k6_mesfunc1 X0 X1 X2))\wedge(m1_subset_1 (k6_mesfunc1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v3_membered X1)\wedge((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 (k32_valued_1 X0 X1 X2))\wedge(m1_subset_1 (k32_valued_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ ((v1_relat_1 (k30_valued_1 X0)) \wedge ((v1_funct_1 (k30_valued_1 \\ X0)) \wedge (v1_valued_0 (k30_valued_1 X0)))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k1_real_1 X0) k1_numbers) \quad (13)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ ((v1_funct_1 (k1_mesfunc5 X0 X1)) \wedge (m1_subset_1 (k1_mesfunc5 \\ X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers)))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ k1_mesfunc5 X0 X1 = X1)) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ (k30_valued_1 X0 = k24_valued_1 X0 (k4_xcmplx_0 np_1)) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \Rightarrow ((v1_relat_1 X0) \wedge (v1_valued_0 X0)) \quad (17)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v5_relat_1 X0 k1_numbers)) \Rightarrow ((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \quad (18)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \quad (19)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (20)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \quad (21)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\\ (r2_relset_1 X0 k7_numbers (k7_mesfunc1 X0 (k1_mesfunc5 X0 X1)) \\ (k1_mesfunc5 X0 (k26_valued_1 X0 k1_numbers X1 (k1_real_1 np_1)))) \wedge \\ (r2_relset_1 X0 k7_numbers (k7_mesfunc1 X0 (k1_mesfunc5 X0 X1)) \\ (k1_mesfunc5 X0 (k32_valued_1 X0 k1_numbers X1)))) \end{aligned}$$