

t28\_relat\_1  
(TMLx8ed7fLWHavYLLZyavxbkXLUUij8EX6E)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_relat\_1 X0) \Rightarrow (\forall X1.(v1\_relat\_1 X1) \Rightarrow (r1\_tarski (k10\_xtuple\_0 (k3\_relat\_1 X0 X1)) (k10\_xtuple\_0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.v1\_relat\_1 (k3\_relat\_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(v1\_relat\_1 X2) \Rightarrow ((X2 = k3\_relat\_1 X0 X1) \Leftrightarrow (\forall X3.\forall X4.(k4\_tarski X3 X4 \in X2) \Leftrightarrow (\exists X5.(k4\_tarski X3 X5 \in X0) \wedge (k4\_tarski X5 X4 \in X1)))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(r1\_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k10\_xtuple\_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.k4\_tarski X3 X2 \in X0)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k9\_xtuple\_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.k4\_tarski X2 X3 \in X0)) \quad (6)$$

**Theorem 1**

$$\forall X0.(v1\_relat\_1 X0) \Rightarrow (\forall X1.(v1\_relat\_1 X1) \Rightarrow ((r1\_tarski (k9\_xtuple\_0 X0) (k10\_xtuple\_0 X1)) \Rightarrow (k10\_xtuple\_0 (k3\_relat\_1 X1 X0) = k10\_xtuple\_0 X0)))$$