

t28_xtuple_0 (TMaaGDYND- MYM2XW75fSYzDnscTixcDLu4KG)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. r1_tarski (k3_xboole_0 X0 X1) X0 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. k4_tarski X3 X2 \in X0)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. r1_tarski (k10_xtuple_0 (k3_xboole_0 X0 X1)) (k3_xboole_0 (k10_xtuple_0 X0) (k10_xtuple_0 X1))$$