

t292_xxreal_1
(TMKY9tBK2WgQ33t91fmirn8uhV8sVMW7yi7)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\neg(X0 \in k1_numbers) \wedge (r1_xxreal_0 k1_xxreal_0 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k6_subset_1 (k3_xxreal_1 X2 X1) (k3_xxreal_1 X0 X1) = k3_xxreal_1 X2 X0)))) \quad (2)$$

Assume the following.

$$v1_xxreal_0 k1_xxreal_0 \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Leftrightarrow (X0 \in k1_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (5)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (k6_subset_1 (k3_xxreal_1 X0 k1_xxreal_0) (k3_xxreal_1 X1 k1_xxreal_0) = k3_xxreal_1 X0 X1))$$