

t29_absvalue

(TMKQoUWqwLfsnD8wfQFUP7CpKhtqq2iJaj9)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (r1_xreal_0 k6_numbers X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k18_complex1 X0 = k16_complex1 X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (((r1_xreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = X0)) \wedge ((\neg r1_xreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = k4_xcmplx_0 X0))) \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (4)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (5)$$

Theorem 1 $\forall X0.(v7_ordinal1 X0) \Rightarrow (X0 = k18_complex1 X0)$.