

t29_bvfunc_8

(TMMDsnJyN45qDauTqF3mzyZY1Xf3k2nn1oh)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_bvfunc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\
 & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((r2_funct_2 X0 k6_margrel1 \\
 & (k9_bvfunc_1 X0 X3 (k9_bvfunc_1 X0 X2 X1)) (k12_bvfunc_1 X0)) \Rightarrow (\\
 & r2_funct_2 X0 k6_margrel1 (k9_bvfunc_1 X0 X2 (k9_bvfunc_1 X0 X3 \\
 & X1)) (k12_bvfunc_1 X0))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\
 & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\
 & (k9_bvfunc_1 X0 (k9_bvfunc_1 X0 X1 (k9_bvfunc_1 X0 X2 X3)) (k9_bvfunc_1 \\
 & X0 (k9_bvfunc_1 X0 X1 X2) (k9_bvfunc_1 X0 X1 X3))) (k12_bvfunc_1 \\
 & X0))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 \\ & X1)\wedge((v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1))))))\wedge((v1_funct_1 X2)\wedge((v1_funct_2 \\ & X2 X0 k6_margrel1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1))))))\Rightarrow((v1_funct_1 (k9_bvfunc_1 X0 X1 X2))\wedge(\\ & (v1_funct_2 (k9_bvfunc_1 X0 X1 X2) X0 k6_margrel1)\wedge(m1_subset_1 \\ & (k9_bvfunc_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(\\ & (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(\forall X3.((v1_funct_1 X3)\wedge \\ & ((v1_funct_2 X3 X0 k6_margrel1)\wedge(m1_subset_1 X3 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(r2_funct_2 X0 k6_margrel1 \\ & (k9_bvfunc_1 X0 (k9_bvfunc_1 X0 X1 X2) (k9_bvfunc_1 X0 (k9_bvfunc_1 \\ & X0 X1 (k9_bvfunc_1 X0 X2 X3)) (k9_bvfunc_1 X0 X1 X3))) (k12_bvfunc_1 \\ & X0)))) \end{aligned}$$