

t29\_complfld (TMd-  
WwF8Lt2EoNwnmZJPCsgPKC16qiDq6cGZ)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_complfld : \iota$  be given. Let  $k3\_vectsp\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k6\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v36\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $l6\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $k6\_complex1 : \iota$  be given. Let  $k2\_numbers : \iota$  be given. Let  $u1\_algstr\_0 : \iota \Rightarrow \iota$  be given. Let  $k27\_binop\_2 : \iota$  be given. Let  $u2\_algstr\_0 : \iota \Rightarrow \iota$  be given. Let  $k29\_binop\_2 : \iota$  be given. Let  $k5\_complex1 : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 k1\_complfld)) \Rightarrow (\forall X1. \\ & (m1\_subset\_1 X1 (u1\_struct\_0 k1\_complfld)) \Rightarrow (\forall X2.(v1\_xcmplx\_0 \\ & X2) \Rightarrow (\forall X3.(v1\_xcmplx\_0 X3) \Rightarrow (((X0 = X2) \wedge (X1 = X3)) \Rightarrow ((X1 = \\ & k4\_struct\_0 k1\_complfld) \vee (k3\_vectsp\_1 k1\_complfld X0 X1 = k6\_binop\_2 \\ & X2 X3)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmplx\_0 X1) \Rightarrow ((k7\_xcmplx\_0 X0 X1 = np\_1) \Rightarrow (X0 = X1))) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k6\_binop\_2 X0 X1 = k7\_xcmplx\_0 X0 X1) \tag{3}$$

Assume the following.

$$(v36\_algstr\_0 k1\_complfld) \wedge (l6\_algstr\_0 k1\_complfld) \tag{4}$$

Assume the following.

$$k6\_complex1 = np\_1 \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.((v36\_algstr\_0 X0) \wedge (l6\_algstr\_0 X0)) \Rightarrow ((X0 = k1\_complfld) \Leftrightarrow \\ ((u1\_struct\_0 X0 = k2\_numbers) \wedge ((u1\_algstr\_0 X0 = k27\_binop\_2) \wedge \\ ((u2\_algstr\_0 X0 = k29\_binop\_2) \wedge ((k5\_struct\_0 X0 = k6\_complex1) \wedge \\ (k4\_struct\_0 X0 = k5\_complex1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 k1\_complfld)) \Rightarrow (v1\_xcmplx\_0 X0) \quad (7)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 k1\_complfld)) \Rightarrow (\forall X1. \\ (m1\_subset\_1 X1 (u1\_struct\_0 k1\_complfld)) \Rightarrow ((k3\_vectsp\_1 k1\_complfld \\ X1 X0 = k5\_struct\_0 k1\_complfld) \Rightarrow ((X0 = k4\_struct\_0 k1\_complfld) \vee \\ (X1 = X0)))) \end{aligned}$$