

t29_euclid

(TMQ3o7u4HUSbKbge9RBUqyJ4hfVjaBGB6MQ)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $k1_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $v5_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge (\\ (v13_algstr_0 X1) \wedge ((v2_rlvect_1 X1) \wedge ((v3_rlvect_1 X1) \wedge ((v4_rlvect_1 \\ X1) \wedge ((v5_rlvect_1 X1) \wedge ((v6_rlvect_1 X1) \wedge ((v7_rlvect_1 X1) \wedge \\ ((v8_rlvect_1 X1) \wedge (l1_rlvect_1 X1)))))))))) \Rightarrow (\forall X2.(m1_subset_1 \\ X2 (u1_struct_0 X1)) \Rightarrow (((X0 = k6_numbers) \vee (X2 = k4_struct_0 X1)) \Rightarrow \\ (k1_rlvect_1 X1 X2 X0 = k4_struct_0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (3)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 \\ X0) \wedge (v1_xreal_0 X0))) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1\ X0) \Rightarrow & ((v2_pre_topc\ (k15_euclid\ X0)) \wedge \\ & ((v13_algstr_0\ (k15_euclid\ X0)) \wedge ((v2_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v3_rlvect_1\ (k15_euclid\ X0)) \wedge ((v4_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v5_rlvect_1\ (k15_euclid\ X0)) \wedge ((v6_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & ((v7_rlvect_1\ (k15_euclid\ X0)) \wedge ((v8_rlvect_1\ (k15_euclid\ X0)) \wedge \\ & (v5_rltopsp1\ (k15_euclid\ X0)))))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow ((\neg v2_struct_0\ (k15_euclid\ X0)) \wedge (v5_rltopsp1\ (k15_euclid\ X0))) \quad (6)$$

Assume the following.

$$\forall X0.(l1_rltopsp1\ X0) \Rightarrow ((l1_rlvect_1\ X0) \wedge (l1_pre_topc\ X0)) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow ((v5_rltopsp1\ (k15_euclid\ X0)) \wedge (l1_rltopsp1\ (k15_euclid\ X0))) \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.((\neg v2_struct_0\ X0) \wedge (l1_rlvect_1\ X0)) \Rightarrow & ((v8_rlvect_1 \\ X0) \Leftrightarrow (\forall X1.(m1_subset_1\ X1\ (u1_struct_0\ X0)) \Rightarrow & (k1_rlvect_1 \\ X0\ X1\ np_1 = X1))) \end{aligned} \quad (9)$$

Theorem 1

$$\begin{aligned} \forall X0.(v7_ordinal1\ X0) \Rightarrow & (\forall X1.(m1_subset_1\ X1\ (u1_struct_0 \\ (k15_euclid\ X0))) \Rightarrow & ((k1_rlvect_1\ (k15_euclid\ X0)\ X1\ np_1 = X1) \wedge \\ (k1_rlvect_1\ (k15_euclid\ X0)\ X1\ k6_numbers = k4_struct_0\ (k15_euclid & \\ X0)))) \end{aligned}$$