

t29_finseq_7

(TMMVuvZNPjr7iVAWnsnQraKFj59XtUNdRik)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k17_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_rfinseq : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((k7_finseq_1 X0 k1_xboole_0 = X0) \wedge (k7_finseq_1 k1_xboole_0 X0 = X0)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow (k2_rfinseq X0 (k3_finseq_1 X1) X1 = k1_xboole_0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m2_finseq_1 X1 X0) \Rightarrow \\ & \quad (\forall X2.(v7_ordinal1 X2) \Rightarrow (\forall X3.(v7_ordinal1 X3) \Rightarrow (\\ & \quad \quad ((r1_xxreal_0 np_1 X2) \wedge (r1_xxreal_0 X3 (k3_finseq_1 X1))) \Rightarrow (\\ (r1_xxreal_0 X3 X2) \vee (k2_finseq_7 X0 X1 X2 X3 = k8_finseq_1 X0 (k8_finseq_1 \\ X0 (k8_finseq_1 X0 (k8_finseq_1 X0 (k17_finseq_1 X0 (k7_nat_d X2 \\ np_1) X1) (k12_finseq_1 X0 (k7_partfun1 X0 X1 X3))) (k17_finseq_1 \\ X0 (k7_nat_d (k7_nat_d X3 X2) np_1) (k2_rfinseq X0 X2 X1))) (k12_finseq_1 \\ X0 (k7_partfun1 X0 X1 X2))) (k2_rfinseq X0 X3 X1))))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X0) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_finseq_1 X1 X0) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (k8_finseq_1 X0 X1 X2 = k7_finseq_1 X1 X2) \quad (10)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (11)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ & (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ X0)))))) \end{aligned} \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X1 X0)\wedge(m1_finseq_1 X2 X0))\Rightarrow(m2_finseq_1 (k8_finseq_1 X0 X1 X2) X0) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(m1_subset_1 (k7_partfun1 X0 X1 X2) X0) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(m1_subset_1 (k7_nat.d X0 X1) k5_numbers) \quad (17)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (18)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X1)\wedge(m1_finseq_1 X2 X0))\Rightarrow(m2_finseq_1 (k2_rfinseq X0 X1 X2) X0) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X1)\wedge(m1_finseq_1 X2 X0))\Rightarrow(m2_finseq_1 (k17_finseq_1 X0 X1 X2) X0) \quad (21)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (23)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (24)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xxreal_0 X0) \quad (25)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m2_finseq_1 X1 X0) \Rightarrow \\ & (\forall X2.(v7_ordinal1 X2) \Rightarrow ((r1_xxreal_0 np_1 X2) \Rightarrow ((r1_xxreal_0 \\ & (k3_finseq_1 X1) X2) \vee (k2_finseq_7 X0 X1 X2 (k3_finseq_1 X1) = k8_finseq_1 \\ & X0 (k8_finseq_1 X0 (k8_finseq_1 X0 (k17_finseq_1 X0 (k7_nat_d X2 \\ & np_1) X1) (k12_finseq_1 X0 (k7_partfun1 X0 X1 (k3_finseq_1 X1)))) \\ & (k17_finseq_1 X0 (k7_nat_d (k7_nat_d (k3_finseq_1 X1) X2) np_1) \\ & (k2_rfinseq X0 X2 X1))) (k12_finseq_1 X0 (k7_partfun1 X0 X1 X2)))))) \end{aligned}$$