

t29_funct_1 (TMKiPCNYpDEnUiXRU-
gocXRMTvK4xkA8szfT)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k3_relat_1 \\ & X2 X1)) \Rightarrow (k1_funct_1 (k3_relat_1 X2 X1) X0 = k1_funct_1 X1 (k1_funct_1 \\ & X2 X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k3_relat_1 \\ & X2 X1)) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (k1_funct_1 X2 X0 \in k9_xtuple_0 X1)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((\\ & v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k3_relat_1 X0 \\ & X1)) \wedge (v1_funct_1 (k3_relat_1 X0 X1))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k3_relat_1 X0 X1) \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v2_funct_1 X0) \Leftrightarrow \\ & (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \wedge ((X2 \in k9_xtuple_0 \\ & X0) \wedge (k1_funct_1 X0 X1 = k1_funct_1 X0 X2)))) \Rightarrow (X1 = X2))) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{6}$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.(X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (7)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((k10_xtuple_0 (k3_relat_1 X1 X0) = k10_xtuple_0 X0) \wedge (v2_funct_1 X0)) \Rightarrow (r1_tarski (k9_xtuple_0 X0) (k10_xtuple_0 X1))))$$