

t29_glib_000

(TMWbFmCvu1jfsCA6FhoNJ3j97CCqGZwNsi)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k20_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_glib_000 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_glib_000 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1_subset_1 X1 (k1_zfmisc_1 X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (k4_subset_1 X0 X1 X2 = k2_xboole_0 X1 X2) \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0))))) \Rightarrow (m1_subset_1 (k19_glib_000 X0 X1) (k1_zfmisc_1 (k7_glib_000 X0))) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0))))) \Rightarrow (m1_subset_1 (k18_glib_000 X0 X1) (k1_zfmisc_1 (k7_glib_000 X0))) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1. k20_glib_000 \\ X0 X1 = k4_subset_1 (k7_glib_000 X0) (k18_glib_000 X0 X1) (k19_glib_000 \\ X0 X1)) \end{aligned} \tag{6}$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1. (r1_tarski \\ (k18_glib_000 X0 X1) (k20_glib_000 X0 X1)) \wedge (r1_tarski (k19_glib_000 \\ X0 X1) (k20_glib_000 X0 X1))) \end{aligned}$$