

## t29\_hilbert2

(TMad2F JrMao7WHDEgYjvGt2pDcQTV5Jo5Kf)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_hilbert1 : \iota$  be given. Let  $k1\_hilbert2 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $np\_3 : \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((X1 = k9\_finseq\_1 X0) \Leftrightarrow ((k3\_finseq\_1 X1 = np\_1) \wedge (k1\_funct\_1 X1 np\_1 = X0))) \quad (1)$$

Assume the following.

$$m1\_subset\_1 k1\_xboole\_0 k4\_ordinal1 \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_3) \wedge (m2\_subset\_1 np\_3 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_3 k5\_numbers) \wedge (m1\_subset\_1 np\_3 k1\_numbers)) \quad (3)$$

Assume the following.

$$\neg v1\_xboole\_0 np\_3 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2. (m2\_subset\_1 X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0.k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (6)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (7)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow(k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (10)$$

Assume the following.

$$\forall X0.v1\_finseq\_1 (k5\_finseq\_1 X0) \quad (11)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (12)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k5\_finseq\_1 X0))\wedge(v1\_funct\_1 (k5\_finseq\_1 X0)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge((\neg v1\_xboole\_0 X1)\wedge(v7\_ordinal1 X1)))\Rightarrow(\neg v1\_xboole\_0 (k2\_xcmplx\_0 X0 X1)) \quad (14)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (15)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(m2\_subset\_1 (k2\_nat\_1 X0 X1) k1\_numbers k5\_numbers) \quad (17)$$

Assume the following.

$$k2\_hilbert1 = k12\_finseq\_1 \ k5\_numbers \ k6\_numbers \quad (18)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 \ X0 \ k5\_numbers) \Rightarrow (k1\_hilbert2 \ X0 = k12\_finseq\_1 \ k5\_numbers \ (k2\_nat\_1 \ np\_3 \ X0)) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 \ X0 \ k5\_numbers) \wedge (v7\_ordinal1 \ X1)) \Rightarrow (k2\_nat\_1 \ X0 \ X1 = k2\_nat\_1 \ X1 \ X0) \quad (20)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 \ X0 \ k4\_ordinal1) \Rightarrow (v7\_ordinal1 \ X0) \quad (21)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 \ X0) \Rightarrow (\forall X1.(m1\_subset\_1 \ X1 \ (k1\_zfmisc\_1 \ X0)) \Rightarrow (v1\_xboole\_0 \ X1)) \quad (22)$$

**Theorem 1**  $\forall X0.(m1\_subset\_1 \ X0 \ k5\_numbers) \Rightarrow (k2\_hilbert1 \neq k1\_hilbert2 \ X0)$ .