

t29_int_2

(TMTmfccxAbGfv5t2cbtbZru43pkWBV2tRrF)

October 27, 2020

Let $v1_int_2 : \iota \Rightarrow o$ be given. Let $np_4 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $r1_int_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & ((v2_xxreal_0\ np_4) \wedge (m2_subset_1\ np_4\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_4\ k5_numbers) \wedge (m1_subset_1\ np_4\ k1_numbers)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0\ np_2) \wedge (m2_subset_1\ np_2\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_2\ k5_numbers) \wedge (m1_subset_1\ np_2\ k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$k3_xcmplx_0\ np_2\ np_2 = np_4 \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0. (v7_ordinal1\ X0) \Rightarrow ((v1_int_2\ X0) \Leftrightarrow ((\neg r1_xxreal_0\ X0 \\ np_1) \wedge (\forall X1. (v7_ordinal1\ X1) \Rightarrow (\neg (r1_int_1\ X1\ X0) \wedge ((X1 \neq \\ np_1) \wedge (X1 \neq X0)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0. (v1_int_1\ X0) \Rightarrow (\forall X1. (v1_int_1\ X1) \Rightarrow ((r1_int_1 \\ X0\ X1) \Leftrightarrow (\exists X2. (v1_int_1\ X2) \wedge (X1 = k3_xcmplx_0\ X0\ X2)))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset.1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_int.1 X0) \quad (8)$$

Theorem 1 $\neg v1_int.2 np_4$.