

t29_lpspace1

(TMS1iBctfDs3UA8PAnyBzXK3GK78dyT9PCe)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r1_lpspace1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow ((r2_relset_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow ((v1_funct_1 (k2_partfun1 X0 X1 X2 X3)) \wedge (m1_subset_1 (k2_partfun1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \quad (2)$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k1_numbers)))) \Rightarrow (\forall X4.((v1_funct_1 X4) \wedge (m1_subset_1 \\
& X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow ((r1_lpspace1 \\
& X0 X1 X2 X3 X4) \Leftrightarrow (\exists X5.(m2_subset_1 X5 (k1_zfmisc_1 X0) X1) \wedge \\
& ((k3_funct_2 X1 k7_numbers X2 X5 = k6_numbers) \wedge (r2_relset_1 X0 \\
& k1_numbers (k2_partfun1 X0 k1_numbers X3 (k3_subset_1 X0 X5)) (\\
& k2_partfun1 X0 k1_numbers X4 (k3_subset_1 X0 X5))))))))) \\
& \tag{3}
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge ((v1_funct_2 \\
& X2 X1 k7_numbers) \wedge ((v10_valued_0 X2) \wedge ((v6_supinf_2 X2) \wedge ((v4_measure1 \\
& X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers)))))) \Rightarrow \\
& (\forall X3.((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 k1_numbers)))) \Rightarrow (\forall X4.((v1_funct_1 X4) \wedge (m1_subset_1 \\
& X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow ((r1_lpspace1 \\
& X0 X1 X2 X3 X4) \Rightarrow (r1_lpspace1 X0 X1 X2 X4 X3))))))
\end{aligned}$$