

t29\_nat\_1 (TMX-  
AxrkVPQp7WRtTHwwwHwaJccecFtE6NEA)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $r1\_xreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_5 : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $np\_3 : \iota$  be given. Let  $np\_4 : \iota$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\neg (r1\_xreal\_0 X0 (k1\_nat\_1 X1 np\_1)) \wedge (\neg r1\_xreal\_0 X0 X1) \wedge (X0 \neq k1\_nat\_1 X1 np\_1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\neg (r1\_xreal\_0 X0 np\_4) \wedge ((X0 \neq k6\_numbers) \wedge ((X0 \neq np\_1) \wedge ((X0 \neq np\_2) \wedge ((X0 \neq np\_3) \wedge (X0 \neq np\_4)))))) \quad (2)$$

Assume the following.

$$((v2\_xreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (3)$$

Assume the following.

$$k2\_xcmplx\_0 np\_4 np\_1 = np\_5 \quad (4)$$

Assume the following.

$$k2\_xcmplx\_0 np\_3 np\_1 = np\_4 \quad (5)$$

Assume the following.

$$k2\_xcmplx\_0 np\_2 np\_1 = np\_3 \quad (6)$$

Assume the following.

$$k2\_xcmplx\_0 np\_1 np\_1 = np\_2 \quad (7)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (8)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(m1\_subset\_1 X1 k5\_numbers))\Rightarrow (k1\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(v7\_ordinal1 X1))\Rightarrow(v7\_ordinal1 (k2\_xcmplx\_0 X0 X1)) \quad (11)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (12)$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow(\neg(r1\_xxreal\_0 X0 np\_5)\wedge((X0\neq k6\_numbers)\wedge((X0\neq np\_1)\wedge((X0\neq np\_2)\wedge((X0\neq np\_3)\wedge((X0\neq np\_4)\wedge(X0\neq np\_5))))))))$$