

t29_setlim_1

(TMbQiCyFg8kFBzqcYncLC7ZGejd7G1jLTpv)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_kurato_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_card_3 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (k1_prob_1 X0 X1 = k3_card_3 \\ X1) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (k1_kurato_0 X0 X1 = k3_card_3 \\ X1) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k3_subset_1 \\ X0 (k3_subset_1 X0 X1) = X1) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (k2_prob_1 X0 (k2_prob_1 X0 \\ X1) = X1) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers \\ (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (k9_setfam_1 X0))))))\Rightarrow((v1_funct_1 (k2_prob_1 X0 \\ X1))\wedge((v1_funct_2 (k2_prob_1 X0 X1) k5_numbers (k9_setfam_1 X0))\wedge \\ (m1_subset_1 (k2_prob_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ (k9_setfam_1 X0)))))) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers \\ (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (k9_setfam_1 X0))))))\Rightarrow(m1_subset_1 (k1_prob_1 X0 \\ X1) (k1_zfmisc_1 X0)) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers \\ (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (k9_setfam_1 X0))))))\Rightarrow(k3_prob_1 X0 X1 = k3_subset_1 \\ X0 (k1_prob_1 X0 (k2_prob_1 X0 X1))) \end{aligned} \tag{7}$$

Theorem 1

$$\begin{aligned} \forall X0.\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers \\ (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers (k9_setfam_1 X0))))))\Rightarrow(k1_kurato_0 X0 X1 = k3_subset_1 \\ X0 (k3_prob_1 X0 (k2_prob_1 X0 X1))) \end{aligned}$$