

t29_xxreal_2 (TM-
Noodaru9Ph2q2CA7aLwUvA2pPJ2ZTbk1m)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xxreal_2 : \iota \Rightarrow \iota$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Let $m1_xxreal_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ (v1_xxreal_0 X2) \Rightarrow ((X0 \in k1_xxreal_1 X1 X2) \Leftrightarrow ((r1_xxreal_0 X1 X0) \wedge \\ (r1_xxreal_0 X0 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (v2_membered (k1_xxreal_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v2_membered X0) \Rightarrow (\forall X1.(m1_xxreal_2 X1 X0) \Rightarrow (v1_xxreal_0 X1)) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ X1 = k1_xxreal_2 X0) \Leftrightarrow ((m1_xxreal_2 X1 X0) \wedge (\forall X2.(m1_xxreal_2 \\ X2 X0) \Rightarrow (r1_xxreal_0 X1 X2)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.(v2_membered X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ m1_xxreal_2 X1 X0) \Leftrightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((X2 \in X0) \Rightarrow (r1_xxreal_0 \\ X2 X1)))))) \end{aligned} \quad (6)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k1_xxreal_2 (k1_xxreal_1 X0 X1) = X1)))$$