

t2_cohsp_1
(TMZV4ZgL7KQ2LRi94r1vH5Kt6ieEs5th8dm)

October 27, 2020

Let $k3_tarSKI : \iota \Rightarrow \iota$ be given. Let $k1_cohsp_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k1_tarSKI : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X1 \in k1_cohsp_1 X0) \Leftrightarrow (\neg(X1 \neq k1_xboole_0) \wedge (\forall X2. \neg(X1 = k1_tarSKI X2) \wedge (X2 \in X0))) \quad (2)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k3_tarSKI X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X2 \in X3) \wedge (X3 \in X0))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarSKI X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (5)$$

Theorem 1 $\forall X0. k3_tarSKI (k1_cohsp_1 X0) = X0.$