

t2_euclid_5
(TMRB9h32BPWjmb8vTadFrGf9vghjt2ujXoR)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid_5 : \iota \Rightarrow \iota$ be given. Let $k4_euclid_5 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_euclid_5 : \iota \Rightarrow \iota$ be given. Let $k3_euclid_5 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k11_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $np_2 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_monoid_0 : \iota \Rightarrow o$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $v5_rltopsp1 : \iota \Rightarrow o$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $l1_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_monoid_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_relat_1 X3) \wedge \\ & ((v1_funct_1 X3) \wedge (v1_finseq_1 X3))) \Rightarrow ((X3 = k11_finseq_1 X0 X1 \\ & X2) \Leftrightarrow ((k3_finseq_1 X3 = np_3) \wedge ((k1_funct_1 X3 np_1 = X0) \wedge ((k1_funct_1 \\ & X3 np_2 = X1) \wedge (k1_funct_1 X3 np_3 = X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_3) \wedge (m2_subset_1 np_3 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_3 k5_numbers) \wedge (m1_subset_1 np_3 k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1_xreal_0 X0) \wedge ((v1_xreal_0 \\ & X1) \wedge (v1_xreal_0 X2))) \Rightarrow (k4_euclid_5 X0 X1 X2 = k11_finseq_1 X0 X1 \\ & X2) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow((v2_monoid_0 (k15_euclid X0))\wedge(v5_rltopsp1 (k15_euclid X0))) \quad (6)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.(l1_rltopsp1 X0)\Rightarrow((l1_rlvect_1 X0)\wedge(l1_pre_topc X0)) \quad (8)$$

Assume the following.

$$\forall X0.(l1_pre_topc X0)\Rightarrow(l1_struct_0 X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_xreal_0 X0)\wedge((v1_xreal_0 X1)\wedge(v1_xreal_0 X2)))\Rightarrow(m1_subset_1 (k4_euclid_5 X0 X1 X2) (u1_struct_0 (k15_euclid np_3))) \quad (10)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow((v5_rltopsp1 (k15_euclid X0))\wedge(l1_rltopsp1 (k15_euclid X0))) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_3)))\Rightarrow(k3_euclid_5 X0 = k1_seq_1 X0 np_3) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_3)))\Rightarrow(k2_euclid_5 X0 = k1_seq_1 X0 np_2) \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_3)))\Rightarrow(k1_euclid_5 X0 = k1_seq_1 X0 np_1) \quad (14)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0)))\Rightarrow(v3_valued_0 X1)) \quad (15)$$

Assume the following.

$$\forall X0.(l1_struct_0 X0) \Rightarrow ((v2_monoid_0 X0) \Rightarrow (v1_monoid_0 X0)) \quad (16)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (v1_finseq_1 X1)) \quad (17)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (18)$$

Assume the following.

$$\forall X0.((v1_monoid_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \quad (19)$$

Assume the following.

$$\forall X0.(v6_membered X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v7_ordinal1 X1)) \quad (20)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow (\forall X2.(m1_subset_1 X2 k1_numbers) \Rightarrow ((k1_euclid_5 (k4_euclid_5 X0 X1 X2) = X0) \wedge ((k2_euclid_5 (k4_euclid_5 X0 X1 X2) = X1) \wedge (k3_euclid_5 (k4_euclid_5 X0 X1 X2) = X2))))))$$