

## t2\_fib\_num2

(TMNU3hbwL1ryX8QaHiqJeeZkq2GjmuDpbah)

October 27, 2020

Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Let  $v1\_abian : \iota \Rightarrow o$  be given. Let  $k3\_power : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_int\_1 X1) \Rightarrow ((\neg v1\_abian X1) \Rightarrow (k3\_power (k4\_xcmplx\_0 X0) X1 = k4\_xcmplx\_0 (k3\_power X0 X1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (k3\_power np\_1 X0 = np\_1) \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_int\_1 X0) \Rightarrow (v1\_xreal\_0 X0) \quad (4)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (5)$$

### Theorem 1

$$\forall X0.((v1\_int\_1 X0) \wedge (\neg v1\_abian X0)) \Rightarrow (k3\_power (k4\_xcmplx\_0 np\_1) X0 = k4\_xcmplx\_0 np\_1)$$