

t2_fraenkel
(TMQK4Ar5wiMN6fo6zX7tzikSnbDS31E8aFm)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 \in k1_funct_2 X0 X1) \Rightarrow ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \quad (1)$$

Assume the following.

$$\forall X0. \neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (X1 \in X0))) \wedge ((v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (v1_xboole_0 X1))) \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. r1_tarski (k1_funct_2 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))$$