

t2_funct_1

(TMHShocdLbBtETkSE6k9AhmUaGdbM7iQcF4)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. (v1_relat_1 X1) \Rightarrow ((X0 = X1) \Leftrightarrow (\forall X2. \forall X3. (k4_tarski X2 X3 \in X0) \Leftrightarrow (k4_tarski X2 X3 \in X1)))) \quad (1)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (k4_tarski X1 X2 \in X0))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k9_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. k4_tarski X2 X3 \in X0)) \quad (3)$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((k9_xtuple_0 X0 = k9_xtuple_0 X1) \wedge (\forall X2. (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 X0 X2 = k1_funct_1 X1 X2))) \Rightarrow (X0 = X1)))$$