

t2_funct_2

(TMbaXjoBJDaxDEyXQggKp9FmujSqqGpi4aa)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow & (((r1_tarski \\ (k9_xtuple_0 X2) X0) \wedge (r1_tarski (k10_xtuple_0 X2) X1)) \Rightarrow & (m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) & \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow & ((v1_funct_1 X0) \wedge \\ ((v1_funct_2 X0 (k9_xtuple_0 X0) (k10_xtuple_0 X0)) \wedge & (m1_subset_1 \\ X0 (k1_zfmisc_1 (k2_zfmisc_1 (k9_xtuple_0 X0) (k10_xtuple_0 X0)))))) & \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (v1_partfun1 X1 X0) \Leftrightarrow (k1_relset_1 X0 X1 = X0) \tag{5}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((v1_partfun1 X2 X0) \Rightarrow (v1_funct_2 X2 X0 X1)) \end{aligned} \tag{7}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski \\ & (k10_xtuple_0 X1) X0) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 (k9_xtuple_0 \\ & X1) X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (k9_xtuple_0 \\ & X1) X0)))))) \end{aligned}$$