

## t2\_gr\_cy\_2 (TMHYuoYmGHaPuozgxyDoK- WrCewfBDKAppgz)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v2\_group\_1 : \iota \Rightarrow o$  be given. Let  $v3\_group\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $l3\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $r1\_struct\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_group\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_domain\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Let  $k5\_group\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_group\_1 X0) \wedge ((v3\_group\_1 \\ X0) \wedge (l3\_algstr\_0 X0)))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\ X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow ((r1\_struct\_0 \\ (k5\_group\_4 X0 (k6\_domain\_1 (u1\_struct\_0 X0) X2)) X1) \Leftrightarrow (\exists X3. \\ (v1\_int\_1 X3) \wedge (X1 = k5\_group\_1 X0 X3 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_group\_1 X0) \wedge ((v3\_group\_1 \\ X0) \wedge (l3\_algstr\_0 X0)))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\ X0)) \Rightarrow (k5\_group\_1 X0 np\_1 X1 = X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_int\_1 X0) \quad (6)$$

**Theorem 1**

$$\forall X0.((\neg v2\_struct\_0 X0) \wedge ((v2\_group\_1 X0) \wedge ((v3\_group\_1 X0) \wedge (l3\_algstr\_0 X0)))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (r1\_struct\_0 (k5\_group\_4 X0 (k6\_domain\_1 (u1\_struct\_0 X0) X1)) X1))$$