

## t2\_graph\_2

(TMHoaxLx7YidbeEBew1qi4q7UJSmhCUweqJ)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_xreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (( \\ & \quad v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow (\forall X2.((v1\_relat\_1 \\ & X2) \wedge ((v1\_funct\_1 X2) \wedge (v1\_finseq\_1 X2))) \Rightarrow ((X2 = k5\_relat\_1 X1 \\ & (k2\_finseq\_1 X0)) \Rightarrow (r1\_xreal\_0 (k3\_finseq\_1 X2) (k3\_finseq\_1 \\ & \quad X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1\_relat\_1 X2) \wedge (v1\_funct\_1 \\ & X2)) \Rightarrow ((X0 \in k9\_xtuple\_0 (k5\_relat\_1 X2 X1)) \Rightarrow (k1\_funct\_1 (k5\_relat\_1 \\ & \quad X2 X1) X0 = k1\_funct\_1 X2 X0)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((X1 \in k1\_relset\_1 k5\_numbers X0) \Leftrightarrow \\ & ((r1\_xreal\_0 np\_1 X1) \wedge (r1\_xreal\_0 X1 (k3\_finseq\_1 X0)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ & \quad X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v4\_relat\_1 X1 X0))\Rightarrow(k1\_rerset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (6)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (7)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (9)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow((v1\_relat\_1 X0)\wedge((v4\_relat\_1 X0 k5\_numbers)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))) \quad (10)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\Rightarrow(v1\_xboole\_0 X1)) \quad (11)$$

**Theorem 1**

$$\begin{aligned} &\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow \\ &\quad (\forall X1.((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))))\Rightarrow(\forall X2.(m2\_subset\_1 X2 k1\_numbers k5\_numbers)\Rightarrow(( \\ &\quad X0 = k5\_relat\_1 X1 (k2\_finseq\_1 X2))\Rightarrow((r1\_xxreal\_0 (k3\_finseq\_1 X0) (k3\_finseq\_1 X1))\wedge(\forall X3.(m2\_subset\_1 X3 k1\_numbers \\ &\quad k5\_numbers)\Rightarrow(((r1\_xxreal\_0 np\_1 X3)\wedge(r1\_xxreal\_0 X3 (k3\_finseq\_1 X0)))\Rightarrow(k1\_funct\_1 X1 X3 = k1\_funct\_1 X0 X3)))))) \end{aligned}$$