

t2\_midsp\_3  
(TMHYNtp5BWZ6nioyZ6nbEGrqtA6fgHTabJG)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 X0) \Rightarrow (\forall X1. (v7\_ordinal1 X1) \Rightarrow ((X0 \in k2\_finseq\_1 X1) \Leftrightarrow ((r1\_xxreal\_0 np\_1 X0) \wedge (r1\_xxreal\_0 X0 X1)))) \quad (2)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 X0) \Rightarrow (\forall X1. (v7\_ordinal1 X1) \Rightarrow (\neg (r1\_xxreal\_0 X0 X1) \wedge (\forall X2. (v7\_ordinal1 X2) \Rightarrow (X1 \neq k2\_xcmplx\_0 X0 X2)))) \quad (3)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 X1)) \Rightarrow (k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 X0) \Leftrightarrow (X0 \in k4\_ordinal1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_nat\_1 X1 X0) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (9)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 k5\_numbers)\Rightarrow(\forall X1.(m1\_subset\_1 X1 k5\_numbers)\Rightarrow(\neg(X0 \in k2\_finseq\_1 X1)\wedge(\forall X2.(m1\_subset\_1 X2 k5\_numbers)\Rightarrow(\forall X3.(m1\_subset\_1 X3 k5\_numbers)\Rightarrow(\neg(X1 = k2\_nat\_1 (k2\_nat\_1 X2 np\_1) X3)\wedge(X0 = k2\_nat\_1 X2 np\_1))))))$$