

## t2\_partfun2

(TMNB3KjdXC77XAu8MJuoZMLsG2bYuRVgvzX)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v5\_relat\_1 X1 X0)) \Rightarrow (k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (m1\_subset\_1 (k1\_relset\_1 X0 X1) (k1\_zfmisc\_1 X0)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v5\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1))) \Rightarrow (\forall X2. (X2 \in k9\_xtuple\_0 X1) \Rightarrow (k7\_partfun1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \quad (5)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. (X1 = k10\_xtuple\_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9\_xtuple\_0 X0) \wedge (X2 = k1\_funct\_1 X0 X3)))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (8)$$

**Theorem 1**

$$\begin{aligned} &\forall X0.\forall X1.(\neg v1\_xboole\_0 X1)\Rightarrow(\forall X2.(\neg v1\_xboole\_0 \\ &X2)\Rightarrow(\forall X3.((v1\_funct\_1 X3)\wedge(m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ &(k2\_zfmisc\_1 X2 X1))))\Rightarrow((X0 \in k2\_relset\_1 X1 X3)\Leftrightarrow(\exists X4.( \\ &m1\_subset\_1 X4 X2)\wedge((X4 \in k1\_relset\_1 X2 X3)\wedge(X0 = k7\_partfun1 X1 \\ &X3 X4)))))) \end{aligned}$$