

t2\_rewrite3  
(TMLKBwayx21ugJ8qTqxe3FkRrjurNzouYXh)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\neg(X0 \neq k1\_xboole\_0) \wedge (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 \\ & X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow (\forall X2.X0 \neq k7\_finseq\_1 X1 (k9\_finseq\_1 \\ & X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ & X1))) \Rightarrow ((X1 = k9\_finseq\_1 X0) \Leftrightarrow ((k3\_finseq\_1 X1 = np\_1) \wedge (k1\_funct\_1 \\ & X1 np\_1 = X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ & (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ & X1)))) \Rightarrow (k3\_finseq\_1 (k7\_finseq\_1 X0 X1) = k2\_nat\_1 (k3\_finseq\_1 \\ & X0) (k3\_finseq\_1 X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge((\neg v1\_xboole\_0 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))))\Rightarrow(\forall X2.(m2\_subset\_1 X2 X0 X1)\Leftrightarrow(m1\_subset\_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0.k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (6)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.v1\_finseq\_1 (k5\_finseq\_1 X0) \quad (9)$$

Assume the following.

$$\neg v1\_finset\_1 k4\_ordinal1 \quad (10)$$

Assume the following.

$$v6\_membered k4\_ordinal1 \quad (11)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k5\_finseq\_1 X0)\wedge(v1\_funct\_1 (k5\_finseq\_1 X0))) \quad (12)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (13)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow(m2\_subset\_1 (k3\_finseq\_1 X0) k1\_numbers k5\_numbers) \quad (14)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\Rightarrow(v1\_xboole\_0 X1)) \quad (15)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0)\Rightarrow(v1\_finset\_1 X0) \quad (16)$$

Assume the following.

$$\forall X0.(v6\_membered X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 X0)\Rightarrow(v7\_ordinal1 X1)) \quad (17)$$

**Theorem 1**

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow \\ (\neg(X0 \neq k1\_xboole\_0) \wedge (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 \\ X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow (\forall X2. \neg(X0 = k7\_finseq\_1 X1 (k9\_finseq\_1 \\ X2)) \wedge (k3\_finseq\_1 X0 = k2\_xcmplx\_0 (k3\_finseq\_1 X1) np\_1)))) \end{aligned}$$