

## t2\_rfunct\_3 (TMRM- sJx52KAzbiL6Qfuv7QtCTPe8jWBpHD5)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k18\_complex1 : \iota \Rightarrow \iota$  be given. Let  $k7\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_rfunct\_3 : \iota \Rightarrow \iota$  be given. Let  $k2\_rfunct\_3 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_0 : \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $k16\_complex1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k4\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k3\_xcmplx\_0 X0 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k2\_xcmplx\_0 X0 k6\_numbers = X0) \quad (3)$$

Assume the following.

$$(m2\_subset\_1 np\_0 k1\_numbers k5\_numbers) \wedge ((m1\_subset\_1 np\_0 k5\_numbers) \wedge (m1\_subset\_1 np\_0 k1\_numbers)) \quad (4)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (5)$$

Assume the following.

$$k4\_xcmplx\_0 np\_0 = np\_0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 X1)) \Rightarrow (k7\_real\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (7)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (8)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k18\_complex1 X0 = k16\_complex1 X0) \quad (9)$$

Assume the following.

$$\exists X0.(v1\_xboole\_0 X0) \wedge ((v1\_xcmplx\_0 X0) \wedge ((v1\_xxreal\_0 X0) \wedge (v1\_xreal\_0 X0))) \quad (10)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 k6\_numbers) \wedge ((\neg r1\_xxreal\_0 k6\_numbers X1) \wedge (r1\_xxreal\_0 k6\_numbers (k3\_xcmplx\_0 X0 X1))))) \quad (11)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 (k4\_xcmplx\_0 X0) (k4\_xcmplx\_0 X1)) \Rightarrow (r1\_xxreal\_0 X1 X0))) \quad (12)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k4\_xcmplx\_0 (k4\_xcmplx\_0 X0) = X0) \quad (13)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow ((v1\_xcmplx\_0 (k4\_xcmplx\_0 X0)) \wedge (v1\_xreal\_0 (k4\_xcmplx\_0 X0))) \quad (14)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (m1\_subset\_1 (k2\_rfunct\_3 X0) k1\_numbers) \quad (15)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (m1\_subset\_1 (k1\_rfunct\_3 X0) k1\_numbers) \quad (16)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (k2\_rfunct\_3 X0 = k4\_xxreal\_0 (k4\_xcmplx\_0 X0) k6\_numbers) \quad (17)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (k1\_rfunct\_3 X0 = k4\_xxreal\_0 X0 k6\_numbers) \quad (18)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (k16\_complex1 X0 = X0)) \wedge ((\neg r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (k16\_complex1 X0 = k4\_xcmplx\_0 X0))) \quad (19)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 X1 X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X0)) \wedge ((\neg r1\_xxreal\_0 X1 X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X1))) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (k4\_xxreal\_0 X0 X1 = k4\_xxreal\_0 X1 X0) \quad (21)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k2\_xcmplx\_0 X0 X1 = k2\_xcmplx\_0 X1 X0) \quad (22)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xxreal\_0 X0) \quad (23)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (24)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (v1\_xreal\_0 X0) \quad (25)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (k18\_complex1 X0 = k7\_real\_1 (k1\_rfunct\_3 X0) (k2\_rfunct\_3 X0))$$