

t2_rlsb_1

(TMV8coNpaQ2FL6CnqPmno5drjhcQoQdda9o)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v1_rlsub_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k3_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (v13_algstr_0 X0) \wedge (v2_rlvect_1 \\ & X0) \wedge (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge (v7_rlvect_1 X0) \wedge (v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\ & X0)))))) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow \\ & (k4_algstr_0 X0 X1 = k1_rlvect_1 X0 X1 (k1_real_1 np_1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (k1_real_1 X0 = k4_xcmplx_0 X0) \quad (3)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k1_real_1 X0) k1_numbers) \quad (4)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge (v13_algstr_0 X0) \wedge (v2_rlvect_1 \\
& X0) \wedge (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v5_rlvect_1 X0) \wedge \\
& ((v6_rlvect_1 X0) \wedge (v7_rlvect_1 X0) \wedge (v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\
& X0)))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 \\
& X0))) \Rightarrow ((v1_rlsub_1 X1 X0) \Leftrightarrow ((\forall X2.(m1_subset_1 X2 (u1_struct_0 \\
& X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow ((X2 \in X1) \wedge \\
& (X3 \in X1)) \Rightarrow (k3_rlvect_1 X0 X2 X3 \in X1)))) \wedge (\forall X2.(m1_subset_1 \\
& X2 k1_numbers) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow \\
& ((X3 \in X1) \Rightarrow (k1_rlvect_1 X0 X3 X2 \in X1))))))
\end{aligned} \tag{5}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge (v13_algstr_0 X0) \wedge (v2_rlvect_1 \\
& X0) \wedge (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v5_rlvect_1 X0) \wedge \\
& ((v6_rlvect_1 X0) \wedge (v7_rlvect_1 X0) \wedge (v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\
& X0)))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 \\
& X0))) \Rightarrow ((v1_rlsub_1 X1 X0) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\
& X0)) \Rightarrow ((X2 \in X1) \Rightarrow (k4_algstr_0 X0 X2 \in X1))))))
\end{aligned}$$